AGRICULTURAL WASTE PROBLEM CAN BE SOLVED WITHIN INDUSTRIAL SYMBIOSIS

G. Filik^{1,*}, Z. Dogan² and A. G. Filik¹

¹Kirsehir Ahi Evran University, Agriculture Faculty, Agricultural Biotechnology Department, 40100, Kirsehir-Turkey; ²Kirsehir Ahi Evran University, Natural and Applied Institute, Agricultural Biotechnology Department, 40100, Kirsehir-Turkey

*Corresponding author's e-mail: gfilik@ahievran.edu.tr

The industrial symbiosis network means the disposal of industrial wastes or their evaluation in different places in order to create a more livable environment and world. In this reason industrial symbiosis network, which is rapidly spreading to the world with the example of Kalundborg-Denmark, has reached significant dimensions both in terms of recycling of the wastes and protecting the ecosystem and bringing them into the economy. The present study was designed to show that businesses that produce agricultural and / or industrial waste can use each other's waste as inputs. In this aim of the current study; 53 enterprises were used in food (16), mining and industry (7), agriculture and livestock (9), energy (6), textile (4) and other (11) sectors. Also, this research has been made to support the establishment of new organized industrial zones according to the industrial symbiosis network for waste management.

Keywords: Industrial symbiosis, industrial ecology, environment, industry, agricultural waste.

INTRODUCTION

The technology developed in parallel with the rapid increase in the world population has made people more consuming society. In order to meet the increasing demand, it is important to present new solutions and to carry on vital activities. The world population is known to be 7.7 billion. This rapid increase threatens the vital space in the world and brings environmental and economic concerns. At this point, innovative approaches that emphasize communication and cooperation are put forward in the new world order where the borders separating countries and sectors are removed. These approaches improve the culture of the sectors and provide added value to the sectors / organizations in the symbiosis network and take care of the environment and social benefit. In this context, the industrial symbiosis, which is the whole system; offering multipurpose opportunities for institutions that have a new understanding and vision, aims to bring together a dynamic network. Industrial symbiosis helps to reduce the environmental impact, provide every participant in the economic structure and develop a culture of cooperation among various industrial organizations. Industrial symbiosis, which refers to the same approach as industrial ecology, aims to establish long-term partnerships improving environmental performance and competitiveness of two or more sectors / organizations which are physically close to each other but normally work independently. This approach is not only to prevent industrial problems but also to provide economic returns to bring together the industrial symbiosis network of different industries and organizations (Ulutas, 2011).

In the second half of the twentieth century, the environmental adversities, caused by the rapidly developing technological and industrial developments and the rapid decline of the nonrenewable natural resources, are increasing in the world. The wastes, generated in parallel with industrialization, have increased significantly over time and the local environmental problems, experienced by these wastes, have gained a global dimension. It is also the part of the sustainable rural and agricultural development (Gülçubuk et al., 2015). Nowadays, industrial symbiosis helps to reduce environmental damages, develops the culture of cooperation between various industrial organizations and provides economic return to every participant. In this context, physical change of materials, energy, water or/and by-products among the organizations; all kinds of resources, logistics and an efficient use of resources and expertise that will allow the company to move in together thinking along the common benefits are realized in other areas (Chertow, 2000). In this respect, the environmental management systems that can be designed for industrial zones such as Organized Industrial Zone (OIZ), by-products, residues or wastes produced by an enterprise can be used as raw material for another enterprise. In this way, industrial problems are not only prevented by environmental problems, but also provide economic return.

Industrial symbiosis encountered in Denmark in the 1970s to the first application in the world. Today, is being applied in many countries such as the UK, Norway, the Netherlands, Australia, Germany, Italy, America, South Korea, China, Mexico, Brazil and Australia. The national industrial symbiosis program in the UK since 2005 is one of the most successful examples in the world (Anonymous, 2012). Within

the program, while the savings of enterprises increased by about £ 250 million per year, 9 million tons of CO₂ emissions decreased by eight million tons. BTC Company in 2009 as the first in Turkey (Baku-Tbilisi-Ceyhan Oil Pipeline Company); In order to support sustainable regional development in the pipeline route, the industrial symbiosis which came into the agenda with the project realized in the Gulf of Iskenderun was carried out after various feasibility and evaluation studies of the company. The implementation phase started in 2011 TTGV (Technology Development Foundation of Turkey) and the application phase of the project with the cooperation agreement between BTC Company began in TTGV executive. The project work, which ended in 2012, was cooperated with International Synergies Limited (ISL), the executive director of the UK National Industrial Symbiosis Program (NISP), and the Environmental Engineering

Department of Middle East Technical University (TTGV, 2015).

Clustering Study: Approximately 1.2 billion tons of petroleum agricultural waste is generated annually worldwide (Anonymous, 2018a). Therefore, the disposal of agricultural waste produced in the world in general or giving back to the economy are very important in terms of environmental cycle. This objective will be made an example of by industrial symbiosis network clustering study recognized that agricultural waste. In current study 53 enterprises were used including food (16), mining and industry (7), agriculture and livestock (9), energy (6), textile (4) and other (11) sectors (Fig. 1). Prepared sectoral based industrial symbiosis samples will help new companies to use waste as inputs to other companies. In addition, companies will have the opportunity to easily manage waste management before the establishment of Organized Industrial Zone (OIZ).




Figure 1. Industrial Symbiosis Network Example.

Table 1. Industrial Symbiosis Network Example.

	<u>le 1. Industrial Symbiosis</u>	Network Example.		
No.	Factory or Plant	Raw Material or Input	Waste or Output	References
1	Cattle Farm	Roughage and Concentrate	Animal Manure, Straw Waste	Öztürk, 2005;
		Feed, Energy		Tolay et al., 2008;
				Kilic et al., 2018
2	Integrated Poultry Plant	Feed, Energy, Packaging	Packaging Waste, Chicken	Görgün et al., 2018;
_	integrated Fourtry Franc	Material	Manure, Bone, Giblets, Blood	Uzun and Sargin, 2018
		Widterful	Flour	Ozun und Surgin, 2010
2	Varminammast Dlant	Household Wests, Comment		Taylor et al. 2002.
3	Vermicompost Plant	Household Waste, Compost,	Vermicompost	Taylor et al., 2003;
		Pulp, Animal Manure	E' LO'LE' LW	Erşahin, 2007
4	Canned Fish Factory	Feed, Energy, Packaging	Fish Oil, Fish Waste,	Kilinc, 2007;
_		Material	Packaging Waste	Akagündüz, 2010
5	Feed Production Plant	Grain, Pulp, Energy, Feed	Packaging Waste, Powder,	Horuz et al., 2015
		Additives	Grain Shell, Grain Waste	
6	Integrated Cotton	Cotton, Energy	Cotton Seed, Ginning, Cotton	Usta et al., 2003;
	Production Factory		Waste	Alkaya, 2010
7	Integrated Tobacco Plant	Tobacco, Energy, Packaging	Tobacco Oil, Waste Water	Çalişkan et al., 2009; Çerçioğlu, 2011
		Material		
8	Pistachio Production Plant	Pistachios, Manure, Energy	Pistachio Glume, Pistachio	Salan and Almab, 2014; Anonymous,
		, , ,	Shell	2018b
9	Integrated Olive Oil Factory	Olive, Glass Bottle, Energy	Olive Mill Waste Water, Olive	Seferoğlu et al., 2008;
		,,,,,	Pomace, Olive Tree Branch	Bain et al., 2010;
			Part	Aktaş and Salih, 2014
10	Tea Production Factory	Tea, Energy, Packaging	Tea Waste, Packaging Waste	Aşik and Kütük, 2012;
10	rea r roduction r actory	Material	Tea waste, I ackaging waste	Bilgin et al., 2016
11	Integrated Orange	Energy, Manure, Packaging	Orange Pulp, Packaging	Günkaya et al., 2016;
11	Production Plant	Material	Waste, Waste Water	
10			*	Filik and Kutlu, 2018
12	Integrated Banana	Energy, Manure, Packaging	Banana Cob, Banana Leaf,	Sözer and Yaldiz, 2011; Topbaşli and
10	Production Plant	Material	Packaging Waste	Sevinçli, 2017
13	Integrated Fruit Juice	Fresh Fruit, Energy,	Fruit Pulp, Waste Water	Yang and Feng, 2008;
	Factory	Packaging Materials		Yalçinkaya et al., 2012;
				Bektaş and Gülmez, 2012
14	Sugar Refinery	Sugar Beet, Energy	Pulp, Cake, Bioethanol, Waste	Kahyaoğlu and Konar, 2006; Dönmez,
			Water	2010;
				Altunbay et al., 2016
15	Flour Production Plant	Wheat, Corn, Energy,	Bran, Razmol, Waste Water,	Fane and Fell, 1977;
		Packaging Materials	Wheat Red-Dog, Wheat Germ	Ertop et al., 2016
16	Cheese Production Facility	Milk, Energy, Yeast,	Whey, Packaging Waste	Sözer and Yaldız, 2006; Dinçoğlu and
	Ž	Packaging Materials	<i>y</i> , <i>c c</i>	Ardıç, 2012
17	Biscuit Production Facility	Whey, Sugar, Oil, Flour	Packaging Waste, Waste Water	
		, ,,,,	88	Khare and Bundela, 2014
18	Jam Production Facility	Fruit, Rose Leaf, Sugar,	Packaging Waste, Waste Water	
10	Jam I roduction I denity	Packaging Materials	i achaging music, music mater	Tallino and Danotti, 2017
19	Integrated Rose Production	0 0	Rose Pulp, Waste Water	Onursal, 2006; Akar, 2012
17	Facility	Rose, Manure, Energy, Packaging Materials	Rose i uip, waste water	Onursai, 2000, Akar, 2012
20	Chips Production Plant		Poteto Apple Wests Wt- O'l	Çanakçi and Özsezen, 2005; Uçaroğlu
20	Chips Production Plant	Potato, Apple, Vegetable	Potato-Apple waste, waste Off	, ,
2.1	T ID . D . I	Oil, Energy	D 1 1 W . D W .	et al., 2016
21	Integrated Beer Production	Barley, Energy, Packaging	Packaging Waste, Beer Yeast	Kaur and Saxena, 2004;
	Plant	Materials		Han et al., 2006;
				Ferreira et al., 2010;
				Bain et al., 2010
22	Integrated Petroleum	Crude Oil, Coal, Energy	Asphalt	Nezahat et al., 2000
	Production Facility			
23	Integrated Coal Production	Coal, Petroleum, Energy,	Coal Powder, Packaging Waste	Aydoğan et al., 1990
	Plant	Packaging Materials	, , ,	
24	Integrated Lime Industry	Limestone, Energy	Lime Waste, Waste Heat	Yang and Feng, 2008;
	Plant	, —	,	Akyarli et al., 2009
25	Integrated Marble Plant	Raw Marble, Energy	Marble Powder, Palette,	Akbulut and Gürer, 2006;
23	mogration martin in I milit	in maiore, Energy	Rubble	Timur and Kiliç, 2013
			Rubbic	rimur and rxing, 2013

015
015
015
3
5
ici, 2017
2006
, 2013
, 2016
06; Çengelci et
16; Torshizi and
O

According to this study wastes of different sectors can be taken as product (input) of other sector and processed. Thus, connections between various companies can be established and can provide environmental health, gain from the field and economic contribution. In the Table 1 it is stated that which sectors inputs (products) and outputs (wastes) can be used in the industrial symbiosis network.

If the industrial network is not used, only some examples of the damage caused to the environment by some of the wastes will be seen more clearly in terms of economy and the environment:

Lime: Lime is produced by the heating calcareous or limestone in nature. So as to make lime from calcareous or limestone, the internal temperature of the incinerator must be above 90-1200°C. Thus, lime is converted to calcium oxide in high degrees. This heat of the exhaust gas to the environment during the conversion is between approximately 25-300°C. During the heat treatment the atmosphere together

with the high temperatures resulting from flue fuel gas, carbon monoxide (CO) and emission of various gases by air spheres under great threat trigger global warming (KİSAD, 2015). For example, 3.5 million years ago, only one kind of natural causes disappeared for a thousand years, while in the 2000s, this figure will increase from 20 thousand to 50 thousand (Anonymous, 2014). Owing to the negative effects of the wastes and fumes of the factories around the Copper Mountains in the USA, the entire plant ecosystem in that region has been destroyed.

Waste water: which is in the negative output of 95% of the industrial enterprises, interferes with the sea and land ecosystem in a variety of ways by interfering with the groundwater, drinking water and seas. Waste water contamination from the contamination of available clean sources puts both humanity and plant and animal populations into the risk zone (TBMM, 2013).

Packaging waste: This will be processed, recycled and collected discarded by invading the area where the earth leads to significant loss of land. Leather, wood, glass, plastic, metal, fabric, slag etc. parts of the wastes are also recycled in terms of environmental health and gain from the area. D-Waste, ISWA, University of Leeds, WtERT, Sweep-Net and SWAPI, which are included in the Waste Atlas Partnership, have produced 2 years of waste atlas. In the report, a map and a profile of the world's 50 largest active waste dumps have been created and it is reported that approximately 40% of the waste produced in the world is disposed of in an unhealthy way in open waste areas. In most of these dumps it is close to urban areas, but also stated that poses a serious threat to human health and the environment. According to this report; waste of 200-300 times the volume of the Giza Pyramid, or 0.6-0.8 km³ of waste, which can affect the daily life of 64 million people (CEVKO, 2015)

As can be seen from the examples given above, industrial symbiosis is the beginning stage of the new generation technology age industry 4.0. In addition, policy makers and local administrators should strive to ensure that investments in their regions are appropriate for the industrial symbiosis network (Chertow, 2007).

Conclusion: In recent years, rapidly evolving consumption habits with industrialization has increased the amount of domestic and industrial waste. Therefore, wastes will be allowed to be evaluated in a way that will not harm the environment. On the one hand, the use of scarce resources and the prevention of environmental pollution on the other hand can be achieved by recycling the wastes to the nature. In particular, rapid and new developments in science and technology have paved the way for the evaluation of industrial wastes. Increasing of new recycling methods and reevaluation of wastes has the potential to prevent environmental pollution. As a result, the spreading of interenterprise cooperation in the structured clustering study is

very important in the evaluation of wastes, which is one of the biggest environmental problems. Through the organized industrial symbiosis network to be used in newly established, new business opportunities will be provided both high rates of economic income.

REFERENCES

- Akagündüz, Ö.Y. 2010. Çipura işleme atıklarının yan ürün olarak değerlendirilmesi. MSc Thesis, Dept. of Fishing Technol. and Seafood Processing Technol., Ege Univ., İzmir, Turkey.
- Akar, E. 2012. Gül posası ile ekolojik tekstil mamul boyama metotları. MSc Thesis, Natural and Applied Institute, Süleyman Demirel Univ., Isparta, Turkey.
- Akbulut, H. and C. Gürer. 2006. Atik mermerlerin asfalt kaplamalarda agrega olarak değerlendirilmesi. İ.M.O Teknik Dergi. 261:3943-3960.
- Aktaş, A. and S. Özer. 2014. Ham Pirina Yağının Biyodizel Potansiyelinin Araştırılması. J. Agr. Fac. of S.D.Ü. 3:132-139.
- Akyarlı, A., A. Kavak, S. Atay and S. Alkaya. 2009. Killi Zeminlerin Kireç İle İyileştirilmesi. TSE Standart Dergisi, Turkey.
- Alkaya, E. 2010. Lif Pamuk Üretimi Yan Ürünlerinin/artıklarının Katma Değerli Ürünlere Dönüştürülmesi: Mevcut Uygulamalar Ve Teknolojik Gelişmeler. II. Ulusal Katı Atık Yönetimi Kongresi, Mersin, Turkey.
- Altunbay, S.G., A. Kangal and S. Gürel. 2016. Bioethanol production from sugar beet. J. Field Crops Cent. Res. Ins. 25:334-339.
- Anonymous. 2012. Türkiye'de Demir-Çelik Sektörü Atıklarının Ekonomiye Kazandırılması. http://www.endustriyelsimbiyoz.org/wp-content/uploads/2013/07/T%C3%BCrkiyede-Demir-%C3%87elik-Sekt%C3%B6r%C3%BCndeki-At%C4%B1klar%C4%B1n-Ekonomiye-Kazand%C4%B1r%C4%B1lmas%C4%B1.pdf Access Date:18.03.15.
- Anonymous. 2014. Haberler. (http://www.bugday.org/portal/haber_detay.php?hid=12 3) Access Date: 18.03.15.
- Anonymous, 2018a. Biyokütle Potansiyeli olarak Tarımsal Atıklar. Biyoenerji. (http://biyoder.org.tr/biyoenerji/biyokutle/tarimsalatiklar/.) Accessed Time: Nov 16 2018.
- Anonymous. 2018b. Bir ton Antep fistiği atığından 14 kilovatsaat elektrik enerjisi üretilecek. (https://enerjienstitusu.org/2014/09/18/bir-ton-antep-fistigi-atigindan-14-kilovatsaat-elektrik-enerjisi-uretilecek/) Accessed Time: Nov. 16. 2018.

- Aşik, B.B. and C. Kütük. 2012. Çay atığı kompostunun çim alanların oluşturulmasında kullanım olanağı. Uludag Üniv. J. Fac Agri. 26:47-58.
- Ateş, Y. 2007. Endüstriyel Atıkların Çevreden İzolasyonunda Kil Kullanımı–Nükleer Atık Örneği. 13. Ulusal Kil Sempozyumu Bildiriler Kitabı, pp.12-14.
- Aydoğan, S., O. Keçekülah and N. Ay. 1990. Kok Tozlarının Biriketleme ile Değerlendirilmesi. Coal Congress, Turkey. pp.463-470.
- Baiano, A. 2014. Recovery of biomolecules from food wastes a review. Molecules 19:14821-14842.
- Bain, A., M. Shenoy, W. Ashton and M. Chertow. 2010. Industrial symbiosis and waste recovery in an Indian industrial area. Resources, Conservation and Recycling 54:1278-1287.
- Bayhan, Y.K. and G.D. Değermenci. 2017. Kozmetik atık sularından fenton prosesiyle organik madde gideriminin ve kinetiğinin incelenmesi. J. Fac. Eng. Archıt. Gazi. 32:181-188.
- Bektaş, Y. and M. Gülmez. 2012. Biyogaz Destekli Yenilenebilir Hibrid Sistemler İle Enerji Üretimi. Przeglad Elektrotechniczny 88:123-134.
- Bilgin, S., A. Koçer, H. Yılmaz, M. Acar and M. Dok. 2016. Çay fabrikası atıklarınının peletlenmesi ve pelet fiziksel özelliklerinin belirlenmesi. J. Agr. Fac. Gaziosmanpasa Univ. 33:70-80.
- Bošnjaković, M. 2013. Wind power buildings integration. (https://bib.irb.hr/datoteka/731699.Bosnjakovic-wind turbine journal.pdf) Access Date:18.03.15.
- Budak, T.B. and E. Çobanoğlu. 2013. Paracetamol İçeren İlaç Fabrikası Atık Sularının Elektrokimyasal Yöntem Kullanılarak Arıtılması. 1. Ilaç Kimyasi, Üretimi, Teknolojisi, Standardizasyonu Kongresi, Kimyagerler Derneği, Antalya, Turkey.
- Chertow, M.R. 2000. Industrial symbiosis: literature and taxonomy. Ann. Rev. Energy Environ. 25:313-337.
- Chertow, M.R. 2007. "Uncovering" Industrial Symbiosis. J. Indust. Ecol. 11:11-30. doi:10.1162/jiec.2007.1110
- Çalışkan, Ö., M.S. Odabaş and N. Çamaş. 2009. Tütün (*Nicotiana tabacum* L.) tohumunun biyoyakıt olarak değerlendirilmesi. J. Agr. Fac. SDÜ 8:46-55.
- Çanakçı, M. and A.N. Özsezen. 2005. Atık yemeklik yağların alternatif dizel yakıt olarak değerlendirilmesi. Gazi Univ. J. Sci. 18:81-91.
- Çengelci, E., H. Bayrakçeken and F. Aksoy. 2011. Hayvansal ve bitkisel yağlardan elde edilen biyodizelin dizel yakıtı ile karşılaştırılması. Electronic J. Vehicle Technol. 3:41-53.
- Çerçioğlu, M. 2011. Sürdürülebilir tarımda tütün atığı kullanım olanakları. J. Agri. Fac. Uludag Univ. 25:101-107.
- ÇEVKO. 2015. Dünyanın En Büyük 50 Çöplüğü. (http://www.cevko.org.tr/index.php?option=com_conten

- t&task=view&id=496&Itemid=229) Access Date:18.03.15
- Çildir, O. and M. Çanakçı. 2006. Çeşitli Bitkisel Yağlardan Biyodizel Üretiminde Katalizör Ve Alkol Miktarının Yakıt Özellikleri Üzerine Etkisinin İncelenmesi. J. Fac. Eng. Archit. Gazi. 21:367-372.
- Demirel, Y. and Y. Çağlar. 2015. Atık Fosfojipslerin Yapı Malzemesi Olarak Ekonomiye Geri Kazandırılması. J. Fac. Eng. Archıt. Gazi. 30:743-750.
- Dinçoğlu, A.H. and M. Ardıç. 2012. Peynir altı suyunun beslenmemizdeki önemi ve kullanım olanakları. Harran Üniv. Vet. Fak. Derg. 1:54-60.
- Dong, L., H. Zhang, T. Fujita, S. Ohnishi, H. Li, M. Fujii and H. Dong. 2013. Environmental and economic gains of industrial symbiosis for Chinese iron/steel industry: Kawasaki's experience and practice in Liuzhou and Jinan. J. Clean. Prod. 59:226-238.
- Dönmez, E.D. 2010. Şeker Fabrikası Atık Sularından Biyogaz Üretimi, Artaş Endüstriyel Teknik Rapor, Turkey. p.29.
- Erşahin, Y.Ş. 2007. Vermikompost ürünlerinin eldesi ve tarımsal üretimde kullanım alternatifleri. JAFAG 24:99-107.
- Ertop, M.H., K. Kutluk, K. Çoşkun and S. Canlı. 2016. Gıda Endüstrisi Yan Ürünleri Kullanımıyla Cips Üretimine Yeni Bir Yaklaşım: Zenginleştirilmiş Gluten Cipsi. Academic Food J. 14:398-406.
- Fane, A.G. and C.J.D. Fell. 1977. Recovery of soluble protein from wheat starch factory effluents. In AIChE Symp. Ser. 73:198-205.
- Ferreira, I.M.P.L.V.O., O. Pinho, E. Vieira and J.G. Tavarela. 2010. Brewer's Saccharomyces yeast biomass: characteristics and potential applications. Trends Food Sci. Technol. 21:77-84.
- Filik, G. and H.R. Kutlu. 2018. Determination of nutrient values in drying citrus pulp with alternative drying methods. BSJ Agri. 1:11-14.
- Görgün, B., E. Bozkurt, D. Kuru, A.A. Borazan and N. Ural. 2018. Atık Tavuk Tüyünün Zemin İyileştirmesinde Değerlendirilmesi. BSEU J. Sci. 5:89-96.
- Gündüzalp, A.A. and S. Güven. 2016. Atık, Çeşitleri, Atık Yönetimi, Geri Dönüşüm ve Tüketici: Çankaya Belediyesi ve Semt Tüketicileri Örneği. Hacettepe Üniversitesi Sosyolojik Araştırmalar E-Dergisi, ISSN. 1304-2823.
- Gülçubuk, B., N. Kızılaslan, D. Eroğlu, M. Kan, E. Soysald and G. Olgun. 2015. Kırsal Kalkınma Paradigması ve Yeni Arayışlar. Türkiye Ziraat Mühendisliği VIII. Teknik Kongresi, Ankara, Turkey.
- Günkaya, Z., R. Demirel and M. Banar. 2016. Portakal kabuğu atıklarından üretilen biyokompozit ambalaj filminin aflatoksinlere karşı etkisinin incelenmesi. Pamukkale Univ. Muh. Bilim Derg. 22:513-519.
- Gür, N. 2014. Soda Sanayi A.Ş.'nin Endüstriyel Simbiyoz Çalışmaları. Erişim:

- http://www.endustriyelsimbiyoz.org/wp-content/uploads/2014/02/soda-sanayinin-endustriyelsimbiyoz-calismalari_2014.02.20.pdf Access Date 30.11.2018
- Han, R., H. Li, Y. Li, J. Zhang, H. Xiao and J. Shi. 2006. Biosorption of copper and lead ions by waste beer yeast. J. Hazard. Mater. 137:1569-1576.
- Horuz, A., A. Korkmaz and G. Akınoğlu. 2015. Biyoyakıt Bitkileri ve Teknolojisi. Toprak Bilimi ve Bitki Besleme Dergisi. 3:69-81.
- Kafle, G.K., S.H. Kim and K.I. Sung. 2013. Ensiling of fish industry waste for biogas production: a lab scale evaluation of biochemical methane potential (BMP) and kinetics. Bioresour. Technol. 127:326-336.
- Kahyaoğlu, M. and V. Konar. 2006. Şeker Fabrikası Atık Maddeleri Kullanılarak Pseudomonas aeruginosa'dan Ramnolipit Biyosürfektanı Elde Edilmesi. Fırat Üniv. Fen ve Müh. Bil. Dergisi. 18:493-498.
- Kalın, V. 2005. Tekstil Atıkları ve Pamuk Linterinden, Kâğıt Hamuru Ve Kâğıt Üretim Koşullarının Belirlenmesi. MSc Thesis, Natural and Applied Institute, Kahraman Maraş Sütçü İmam Univ., Kahramanmaraş, Turkey.
- Kaur, V.I and P.K. Saxena. 2004. Incorporation of brewery waste in supplementary feed and its impact on growth in some carps. Bioresour. Technol. 91:01-104.
- Kavas, T., M.Y. Çelik and A. Evcin. 2004. Cam atıklarının çimento üretiminde katkı maddesi olarak kullanılabilirliğinin araştırılması. 5. Endüstriyel Hammaddeler Sempozyumu, , İzmir, Turkey. pp.114-119.
- Khare, A.P. and H. Bundela. 2014. Applicability of single chamber microbial fuel cell for the electricity generation using waste water obtained from biscuit factory and potato processing factory. Ijesrt. 3:1-5.
- Kılıç, İ., H. Delice, S. Sofu and B. Yıldız. 2018. Hayvancılık İşletmelerinde Endüstriyel Simbiyoz Uygulaması: Bursa Örneği. J. Agr. Fac. of S.D.Ü.10:240-248.
- Kılınç, B. 2007. Balık atıklarının değerlendirilmesi. EgeJFAS 24:315-319.
- KİSAD. 2015. Uyum Sürecinin Sağlıklı Planlanması Bağlamında Avrupa Birliği'nde Kireç Sektörü. (http://anadolukirec.com/arsiv.htm) Access Date:18.03.15
- Kozak, M. 2010. Tekstil atıkların yapı malzemesi olarak kullanım alanlarının araştırılması. Yapı Teknolojileri Elektronik Dergisi 6:62-70.
- Mekhilef, S., R. Saidur and A. Safari. 2011. A review on solar energy use in industries. Renew Sust Energ Rev. 15:1777-1790.
- Murathan, A., A. Asan. and T.A. Abdulkarem. 2013. Çevresel Atıkların Yapı Malzemesinde Değerlendirilmesi. J. Fac. Eng. Archit. Gazi. 28: 409-416.
- Narlıoğlu, N., T. Salan, N.S. Çetin and M.H. Alma. 2018. Evaluation of furniture industry wastes in polymer

- composite production. Furniture and Wooden Material Research Journal 1:78-85.
- Ediz, N., H. Özdağ and M. Seyfettinoğlu. 2000. Petrol Rafineri Atığının Bor Süspansiyonlarında Çöktürücü Olarak Kullanımı. Scientific Mining J. 39:27-38.
- Onursal, E. 2006. Gül işleme atıklarının kompostlaştırılmasında optimum C/N oranlarının belirlenmesi üzerine bir araştırma, MSc Thesis, Natural and Applied Institute, Süleyman Demirel Univ., Isparta, Turkey.
- Öğünç, A.V., A.Ö. Şehirli, B.K. Laçin, F. Ercan, H. Güçlü and G. Topçu. 2016. Sıçanlarda Deneysel Mide Ülseri Modelinde Yeşil Çay Ekstraktı ve Peynir Altı Suyu Proteinlerinin Etkileri. Academic Food J. 14:111-122.
- Özdamar, A. 2000. Dünya ve Türkiye'de Rüzgâr Enerjisinden Yararlanılması Üzerine Bir Araştırma. Pamukkale Univ. Muh. Bilim Derg. 6:133-145.
- Özkan, Ö. 2007. Atık cam ve yüksek firin cüruflu katkılı harçların özellikleri. J. Fac. Eng. Archit. Gazi. 27: 87-94.
- Öztürk, M. 2005. Hayvan gübresinden biyogaz üretimi. Çevre ve Orman Bakanlığı, 5, 8-18, Ankara, Turkey.
- Parıltı, N.B. 2011. Petrokimya Endüstrisi Atıksularının Güneş Işığı İle Oksidasyonu. D.E.Ü Müh. Fak.Müh. Bil. Derg. 13:39-51.
- Pearce, J.M. 2008. Industrial symbiosis of very large-scale photovoltaic manufacturing. Renew. Energ. 33:1101-
- Ruffino, B. and M. Zanetti. 2017. Present and future solutions of waste management in a candied fruit–jam factory: Optimized anaerobic digestion for on site energy production. J. Clean Prod. 159:26-37.
- Salan, T. and M.H. Almab. 2014. Antep Fıstığı Atık Kabuklarının Enerji, Kimyasal Madde ve Biyomalzeme Üretiminde Değerlendirilmesinde Kullanılabilecek Termokimyasal Yöntemlere Genel Bir Bakış. Yeşil Altın Antepfıstığı Zirvesi, Yeşil Altın Antepfıstığı Çalıştayı, Gaziantep, Turkey.
- Seferoğlu, S., H.G. Seferoğlu and M.A. Kaptan. 2008. Zeytinyağı fabrikası atığı karasuyun (sıvı) gübre olarak mandarinlere uygulanmasının besin maddesi içeriğine ve meyve kalitesi üzerine etkisi. In 4th Nutrition and Fertilizer Congress. pp.834-840.
- Sözer, S. and O. Yaldız. 2006. Sığır Gübresi ve Peynir Altı Suyu Karışımlarından Biyogaz Üretimi Üzerine Bir Araştırma. Akdeniz Univ. Ziraat Fak. Derg. 19:179-183.
- Sözer, S. and O. Yaldız. 2011. Muz serası atıkları ve siğir gübresi karışımlarından mezofilik fermantasyon sonucu üretilebilecek biyogaz miktarının belirlenmesi üzerine bir araştırma. Akdeniz Univ. Ziraat Fak. Derg. 24:75-78.
- Taşkın, O and A., Vardar. 2016. Tarımsal Üretimde Bazı Yenilenebilir Enerji Kaynakları Kullanımı. Uludag Üniv. J. Fac Agri. 30:179-184.

- Taylor, M., W.P. Clarke and P.F. Greenfield. 2003. The treatment of domestic wastewater using small-scale vermicompost filter beds. Ecol. Eng. 21:197-203.
- TBMM. 2013. Meclis Araştırması Komisyonu Raporu.
 - (http://www.tbmm.gov.tr/sirasayi/donem23/yil01/ss352.pdf) Access Date: 18.03.15
- Timur, M. and H. Kılıç. 2013. Mermer Atıkları Kullanılarak Üretilen Otomotiv Fren Balatasının Sürtünme Katsayısının Farklı Fren Balataları İle Karşılaştırılması. Pamukkale Univ. Muh. Bilim Derg. 19:10-14.
- Tolay, M., H. Yamankaradeniz, S. Yardımcı and R. Reiter. 2008. Hayvansal Atıklardan Biyogaz Üretimi. VII. Ulusal Temiz Enerji Sempozyumu, (UTES), İstanbul, Turkey.
- Topbaşlı, B. and Y. Sevinçli. 2017. Lavanta Sapı Ve Muz Kabuğu Kullanılarak Üretilen Yonga Levhaların Karşılaştırılması. EJOVOC 7:47-53.
- Torshizi, M.V. and A.H. Mighani. 2017. The application of solar energy in agricultural systems. Renewable Energy and Sustainable Development 3:234-240.
- TTGV. 2015. Endüstriyel Simbiyoz Nedir? (https://www.endustriyelsimbiyoz.org/endustriyelsimbiyoz-ve-uygulama-ornekleri/endustriyel-simbiyoz-nedir/) Access Date: 18.03.15
- Uçaroğlu, S., U. Alkan and A. Çataltaş. 2016. Arıtma Çamuru Ve Gübre Karışımının Kompostlanmasına Patates Kabuğu ilavesinin Etkisi. Çevre Bilim ve Teknoloji. 1:134-145.
- Uğurlu, M., A. Gürses and M. Yalçın. 2003. Uçucu kül kullanılarak bazı inorganik iyonların kağıt atık sularından giderimi. Fırat Üniv. Fen ve Müh. Bil. Derg. 15:223-231.
- Ulutaş, F. 2011. Endüstriyel Ekoloji. (http://www.rec.org.tr/dyn_files/20/5923-VI-ENDUSTRIYEL-EKOLOJI.pdf) Access Date: 04.01.18
- Ulutaş, F. 2013. Çimento Sektöründe Endüstriyel Simbiyoz ve Enerji Verimliliği. (http://www.endustriyelsimbiyoz.org/wp-content/uploads/2013/12/%C3%87imento-Se

- kt%C3%B6r%C3%BCnde-End%C3%BCstriyel-Simbiyoz-ve-Enerji-Verimlili%C4% 9Fi_CemenT%C3%BCrk_01.12.2013.pdf) Access Date: 04.01.18
- Usta, I., M. Akalın, D. Kocak, N. Merdan and M. Taşdemir. 2003. Effects of varies quantities of waste cotton fibre in a PP polymer on the mechanical properties of resultant composite structure. GUJS 16:405-410.
- Uysal, F.F. and S. Bahar. 2018. Cüruf Çeşitleri ve Kullanım Alanları. Trakya Univ. J. Eng. Sci. 19:37-52.
- Uzun, M. and S. Sargın. 2018. Tavukçuluk Atıkları Kullanılarak Tekstil Yapı Tasarımı ve Termal Konfor Özelliklerinin İncelenmesi. J TEXT Engin. 25:113-120.
- Üçgül, İ. and B. Turak. 2015. Tekstil Katı Atıklarının Geri Dönüşümü ve Yalıtım Malzemesi Olarak Değerlendirilmesi. Akademik Platform. pp.39-48.
- Ünal, M. and A.V. Katkat. 2003. Bisküvi ve Şekerleme Sanayii Arıtma Çamurunun Toprak Özelliklerine ve Mısır Bitkisinin Kimi Mineral Madde İçeriği Üzerine Etkileri. Ulud. Üniv. Zir. Fak. Derg. 17:107-118.
- Yalçınkaya, M.Y., E. Baytok and M.A. Yörük. 2012. Değişik Meyve Posası Silajlarının Bazı Fiziksel ve Kimyasal Özellikleri. J. Fac. Vet. Med. Univ. Erciyes 9:95-106.
- Yaman, K. 2012. Bitkisel atıkların değerlendirilmesi ve ekonomik önemi. Kastamonu Üniversitesi Orman Fakültesi Dergisi 12:339-348.
- Yang, S. and N. Feng. 2008. A case study of industrial symbiosis: Nanning Sugar Co., Ltd. in China. Resour Conserv Recycl. 52:813-820.
- Yapijakis, C. and L.K. Wang. 2006. Treatment of soap and detergent industry wastes. Waste Treatment in the Process Industries 307.
- Yılmaz, O. 2005. Deri sanayii katı atıklarından aktif karbon üretilebilirliğinin incelenmesi MSc Thesis, Natural and Applied Institute, Ege Univ., İzmir, Turkey.
- Zhu, Q., E.A. Lowe, Y.A. Wei and D. Barnes. 2007. Industrial symbiosis in China: a case study of the Guitang Group. J. Ind. Ecol. 11:31-42.